

Addendum B1

Hepatitis B Vaccine Declination

I understand that due to my occupational exposure to blood or other potentially infectious materials I may be at risk of acquiring hepatitis B virus (HBV) infection. I have been given the opportunity to be vaccinated with hepatitis B vaccine, at my own expense. However, I decline Hepatitis B vaccination at this time. I understand that by declining this vaccine, I continue to be at risk of acquiring hepatitis B, a serious disease. If in the future I continue to have occupational exposure to blood or other potentially infectious materials and I want to be vaccinated with hepatitis B vaccine, I can initiate the vaccination series.

Signature of Student

Date

MMR Vaccine Declination

Adults born after 1957 should receive at least one dose of measles vaccine unless they have already had measles and are immune. (This vaccine can also be given as measles mumps rubella (MMR) vaccine or measles rubella (MR) vaccine.) Those at increased risk of getting measles — college students, international travelers and health care workers — should receive two doses, provided they are given no less than 1 month apart.

More than 95% of the people who receive a single dose of MMR will develop immunity to all three viruses. A second gives immunity to almost all of those who did not respond to the first dose.

Note: *Pregnant women should not receive the MMR vaccine. Also, pregnancy should be avoided 1 month following the receipt of the measles vaccine and 3 months following the MMR vaccine.*

I understand that I may be at risk of contracting a measles, mumps or rubella infection due to my occupational exposure. I have been given the opportunity to be vaccinated with the MMR vaccine, at my own expense. However, I decline the vaccination at this time. I understand that by declining this vaccine, I continue to be at risk of acquiring these infections.

Signature of Student

Date